

233/3

Paper 3

CHEMISTRY – (Practical)



Dec. 2022 – 2¼ hours

Name Index Number

Candidate's Signature Date

482

Instructions to candidates

- (a) Write your name and index number in the spaces provided above.
- (b) Sign and write the date of examination in the spaces provided above.
- (c) Answer **all** the questions in the spaces provided in the question paper.
- (d) You are **not** allowed to start working with the apparatus for the first 15 minutes of the 2¼ hours allowed for this paper. This time is to enable you to read the question paper and make sure you have all the chemicals and apparatus that you may need.
- (e) All working **must** be clearly shown where necessary.
- (f) KNEC mathematical tables and silent electronic calculators may be used.
- (g) **This paper consists of 8 printed pages.**
- (h) **Candidates should check the question paper to ascertain that all the pages are printed as indicated and that no questions are missing.**
- (i) **Candidates should answer the questions in English.**

For Examiner's Use Only

Question	Maximum Score	Candidate's Score
1	20	
2	08	
3	12	
Total Score	40	

0347



317092



Turn over

1. (a) You are provided with the following:

- **Solution A** – Indicator solution
- **Solution B** – 0.05 M compound B
- **Solution C1** – Hydrochloric acid to be used in Questions 1(a) and 1(b)

You are required to determine the concentration in moles per litre of hydrochloric acid in solution C1.

PROCEDURE I (a)

- (i) Place two test tubes in a test tube rack. To the first test tube, place about 2 cm³ of solution B. To the second test tube, place about 2 cm³ of solution C1.
- (ii) Add 2 drops of indicator solution A to each of the test tubes, shake and note the colour of each solution. Record the colours in **Table 1**.

Table 1

Solution	Colour
Solution B + indicator solution A	
Solution C1 + indicator solution A	

(1 mark)

Complete the following statement:

In the titration of solution B (in a conical flask) with hydrochloric acid using indicator solution A, the colour change at the end point is from to
(1 mark)

PROCEDURE II (a)

- (i) Using a pipette and pipette filler, pipette 25.0 cm³ of **solution C1** into a 250 ml volumetric flask. Add distilled water to the mark. Label this as **solution C2**.
- (ii) Fill a burette with **solution C2**.
- (iii) Using a clean pipette and pipette filler, place 25.0 cm³ of **solution B** in a 250 ml conical flask.
- (iv) Titrate **solution B** with **solution C2** using 3 drops of indicator **solution A**. Record the results in **Table 2**.

Table 2

	I	II	III
Final burette reading			
Initial burette reading			
Volume of solution C2 used, cm ³			

(4 marks)

Calculate the:

(i) average volume of **solution C2** used.

(1 mark)

.....

(ii) number of moles of **compound B** used.

(1 mark)

.....

(iii) number of moles of hydrochloric acid used (1 mole of compound B reacts with 2 moles of hydrochloric acid).

(1 mark)

.....

(iv) concentration in moles per litre, of hydrochloric acid in **solution C2**.

(1 mark)

.....

(v) concentration in moles per litre, of hydrochloric acid in **solution C1**.

(1 mark)

.....

- (b) You are provided with two portions of solid D and sodium hydrogen carbonate each weighing 2.5 g.

You are required to determine the heat of reaction of hydrochloric acid with aqueous sodium hydrogen carbonate.

PROCEDURE I (b)

- (i) Using a 100 ml measuring cylinder, measure 30 cm³ of distilled water and place it in a 100 ml plastic beaker.
- (ii) Measure the temperature of the distilled water and record in Table 3.
- (iii) Add one of the portions of solid D to the water. Stir with the thermometer and measure the minimum temperature reached. Record the reading in Table 3.

Table 3

Final temperature of the solution, °C	
Initial temperature of water, °C	
Temperature change, °C	

(1½ marks)

Calculate the:

- (i) heat change of the solution (assume specific heat capacity of solution = 4.2 Jg⁻¹ per degree, density of solution = 1.00g cm⁻³) (1 mark)

.....

.....

.....

.....

- (ii) number of moles of sodium hydrogen carbonate, solid D used (relative formula mass = 84) (1 mark)

.....

.....

.....

- (iii) heat change, ΔH_1 in kJmol^{-1} of sodium hydrogen carbonate (1 mark)

.....

PROCEDURE II (b)

- (i) Clean the 100 ml plastic beaker.
 (ii) Repeat **procedure I (b)** using the second portion of **solid D** and 30 cm^3 of **solution C1** instead of 30 cm^3 of distilled water.
 (iii) Record the results in **Table 4**.

482

Table 4

Final temperature of solution $^{\circ}\text{C}$	
Initial temperature of solution C1, $^{\circ}\text{C}$	
Temperature change, $^{\circ}\text{C}$	

(1½ marks)

Calculate the:

- (i) heat change of the solution (assume specific heat capacity of solution = 4.2 Jg^{-1} per degree, density of solution = 1.00 g cm^{-3}) (1 mark)

.....

- (ii) heat change, ΔH_2 in kJmol^{-1} of sodium hydrogen carbonate (1 mark)

.....

- (iii) heat change, $\Delta H_3 = \Delta H_2 - \Delta H_1$ for the reaction of hydrochloric acid and one mole of aqueous sodium hydrogen carbonate (1 mark)

.....



0347

2. You are provided with an organic compound, **solid M**.

Carry out the following tests and record the observations and inferences in the spaces provided.

- (a) Place about one-fifth of **solid M** on a metallic spatula and burn it using a Bunsen burner flame.

Observations	Inferences

(1 mark)

(1 mark)

- (b) Place the remaining amount of **solid M** in a boiling tube. Add about 15 cm³ of distilled water and shake to dissolve. Use about 2 cm³ portions of the solution, in a test tube, for each of the following tests.

- (i) To the first portion, add 3 drops of acidified potassium dichromate(VI). Warm the mixture.

Observations	Inferences

(1 mark)

(1 mark)

- (ii) To the second portion, add 3 drops of bromine water.

Observations	Inferences

(1 mark)

(1 mark)

- (iii) To the third portion, add **all** the solid sodium carbonate provided. Test any gases produced with a burning splint.

Observations	Inferences

(1 mark)

(1 mark)

3. You are provided with **solution N**. Carry out the following tests and record the observations and inferences in the spaces provided. Use about 2 cm³ portions, in a test tube, for each of the tests.

- (a) To the first portion, add aqueous sodium hydroxide dropwise until in excess.

Observations	Inferences

(1 mark)

(1 mark)

- (b) Warm the second portion and then add aqueous ammonia dropwise until in excess.

Observations	Inferences

(1 mark)

(1 mark)

- (c) To the third portion, add 3 drops of aqueous barium nitrate. Shake and then add about 1 cm³ dilute nitric(V) acid.

Observations	Inferences

(2 marks)

(1 mark)